


Nizal Chandrakumar 10/584,440

=> d his nofile 11-12; d que stat 12; d his nofile 13-14; d que stat 14; d his nofile 15-

FILE 'CASREACT' ENTERED AT 11:31:04 ON 27 DEC 2007
ACT CHANDRAKUMAR/A

L1 STR
L2 3 SEA SSS FUL L1 (10 REACTIONS)

L1 STR



VAR G1=NO2/20/21
VPA 18-11/13 U
VPA 19-1/2/6 U
NODE ATTRIBUTES:
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
RSPEC I
NUMBER OF NODES IS 22

STEREO ATTRIBUTES: NONE
L2 3 SEA FILE=CASREACT SSS FUL L1 (10 REACTIONS)


100.0% DONE 2232 VERIFIED 10 HIT RXNS 3 DOCS
SEARCH TIME: 00.00.02

(FILE 'CASREACT' ENTERED AT 11:31:04 ON 27 DEC 2007)

FILE 'REGISTRY' ENTERED AT 11:31:11 ON 27 DEC 2007
ACT CHANDRAREG/A

L3 STR
L4 36 SEA SSS FUL L3

L3 STR



VAR G1=NO2/20/21
 VPA 18-11/13 U
 VPA 19-1/2/6 U
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RSPEC I
 NUMBER OF NODES IS 22

STEREO ATTRIBUTES: NONE
 L4 36 SEA FILE=REGISTRY SSS FUL L3

100.0% PROCESSED 4782 ITERATIONS 36 ANSWERS
 SEARCH TIME: 00.00.01

FILE 'CAPLUS' ENTERED AT 11:31:20 ON 27 DEC 2007

L5 33 SEA ABB=ON PLU=ON L4
 L6 28 SEA ABB=ON PLU=ON L4/P OR L4 (L) (PREPN/OBI OR PREP/RL)
 L7 72233 SEA ABB=ON PLU=ON HALOGEN?/OBI
 L8 5521 SEA ABB=ON PLU=ON DEALKY?/OBI
 L9 1 SEA ABB=ON PLU=ON L6 AND L8 AND L7
 L10 4 SEA ABB=ON PLU=ON L6 AND (L7 OR L8)
 L11 81104 SEA ABB=ON PLU=ON ETHER#/OBI (L) (REACT?/OBI OR RACT/RL)
 L12 9820 SEA ABB=ON PLU=ON FRIEDEL CRAFT#/OBI
 L13 2 SEA ABB=ON PLU=ON L6 AND (L11 OR L12)
 L14 5 SEA ABB=ON PLU=ON L13 OR L10
 D SCAN TI
 L15 53655 SEA ABB=ON PLU=ON ACYLAT?/OBI
 L16 3 SEA ABB=ON PLU=ON L15 AND L6
 L17 6 SEA ABB=ON PLU=ON L16 OR L14
 L18 6 SEA ABB=ON PLU=ON L9 OR L14 OR L16

FILE 'CASREACT, CAPLUS' ENTERED AT 11:39:28 ON 27 DEC 2007

L19 6 DUP REM L2 L18 (3 DUPLICATES REMOVED)
 ANSWERS '1-3' FROM FILE CASREACT
 ANSWERS '4-6' FROM FILE CAPLUS
 E S SGIYTTETEN A?/AU
 L20 0 SEA ABB=ON PLU=ON SHOUTTEETEN A?/AU
 L21 5 SEA ABB=ON PLU=ON BLEGER F?/AU
 L22 4 SEA ABB=ON PLU=ON MORDACQ F?/AU
 L23 69 SEA ABB=ON PLU=ON PIRON J?/AU


Nizal Chandrakumar 10/584,440

L24 45 SEA ABB=ON PLU=ON SCHOUTEETEN A?/AU
 L25 114 SEA ABB=ON PLU=ON (L20 OR L21 OR L22 OR L23 OR L24)
 L26 2 SEA ABB=ON PLU=ON L25 AND L5
 L27 1 SEA ABB=ON PLU=ON L26 NOT L19

=> fil casreact caplus
 FILE 'CASREACT' ENTERED AT 11:44:25 ON 27 DEC 2007
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'CAPLUS' ENTERED AT 11:44:25 ON 27 DEC 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)


=> d que stat l19
 L1 STR



VAR G1=NO2/20/21
 VPA 18-11/13 U
 VPA 19-1/2/6 U
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ELEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RSPEC I
 NUMBER OF NODES IS 22

STEREO ATTRIBUTES: NONE
 L2 3 SEA FILE=CASREACT SSS FUL L1 (10 REACTIONS)
 L3 STR



VAR G1=NO2/20/21

VPA 18-11/13 U
 VPA 19-1/2/6 U
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RSPEC I
 NUMBER OF NODES IS 22

STEREO ATTRIBUTES: NONE

L4 36 SEA FILE=REGISTRY SSS FUL L3
 L6 28 SEA FILE=CAPLUS ABB=ON PLU=ON L4/P OR L4 (L) (PREPN/OBI OR
 PREP/RL)
 L7 72233 SEA FILE=CAPLUS ABB=ON PLU=ON HALOGEN?/OBI
 L8 5521 SEA FILE=CAPLUS ABB=ON PLU=ON DEALKY?/OBI
 L9 1 SEA FILE=CAPLUS ABB=ON PLU=ON L6 AND L8 AND L7
 L10 4 SEA FILE=CAPLUS ABB=ON PLU=ON L6 AND (L7 OR L8)
 L11 81104 SEA FILE=CAPLUS ABB=ON PLU=ON ETHER#/OBI (L) (REACT?/OBI OR
 RACT/RL)
 L12 9820 SEA FILE=CAPLUS ABB=ON PLU=ON FRIEDEL CRAFT#/OBI
 L13 2 SEA FILE=CAPLUS ABB=ON PLU=ON L6 AND (L11 OR L12)
 L14 5 SEA FILE=CAPLUS ABB=ON PLU=ON L13 OR L10
 L15 53655 SEA FILE=CAPLUS ABB=ON PLU=ON ACYLAT?/OBI
 L16 3 SEA FILE=CAPLUS ABB=ON PLU=ON L15 AND L6
 L18 6 SEA FILE=CAPLUS ABB=ON PLU=ON L9 OR L14 OR L16
 L19 6 DUP REM L2 L18 (3 DUPLICATES REMOVED)

=> d que nos 127

L1 STR
 L2 3 SEA FILE=CASREACT SSS FUL L1 (10 REACTIONS)
 L3 STR
 L4 36 SEA FILE=REGISTRY SSS FUL L3
 L5 33 SEA FILE=CAPLUS ABB=ON PLU=ON L4
 L6 28 SEA FILE=CAPLUS ABB=ON PLU=ON L4/P OR L4 (L) (PREPN/OBI OR
 PREP/RL)
 L7 72233 SEA FILE=CAPLUS ABB=ON PLU=ON HALOGEN?/OBI
 L8 5521 SEA FILE=CAPLUS ABB=ON PLU=ON DEALKY?/OBI
 L9 1 SEA FILE=CAPLUS ABB=ON PLU=ON L6 AND L8 AND L7
 L10 4 SEA FILE=CAPLUS ABB=ON PLU=ON L6 AND (L7 OR L8)
 L11 81104 SEA FILE=CAPLUS ABB=ON PLU=ON ETHER#/OBI (L) (REACT?/OBI OR
 RACT/RL)
 L12 9820 SEA FILE=CAPLUS ABB=ON PLU=ON FRIEDEL CRAFT#/OBI
 L13 2 SEA FILE=CAPLUS ABB=ON PLU=ON L6 AND (L11 OR L12)
 L14 5 SEA FILE=CAPLUS ABB=ON PLU=ON L13 OR L10
 L15 53655 SEA FILE=CAPLUS ABB=ON PLU=ON ACYLAT?/OBI
 L16 3 SEA FILE=CAPLUS ABB=ON PLU=ON L15 AND L6
 L18 6 SEA FILE=CAPLUS ABB=ON PLU=ON L9 OR L14 OR L16
 L19 6 DUP REM L2 L18 (3 DUPLICATES REMOVED)
 L20 0 SEA SHOUTTEETEN A?/AU
 L21 5 SEA BLEGER F?/AU
 L22 4 SEA MORDACQ F?/AU
 L23 69 SEA PIRON J?/AU
 L24 45 SEA SCHOUTEETEN A?/AU
 L25 114 SEA (L20 OR L21 OR L22 OR L23 OR L24)
 L26 2 SEA L25 AND L5
 L27 1 SEA L26 NOT L19

=> d ibib ab fhlt 119 1-3; d .ca hitstr 119 4-6; d .ca 127 1

L19 ANSWER 1 OF 6 CASREACT COPYRIGHT 2007 ACS on STN DUPLICATE 1

ACCESSION NUMBER: 144:450603 CASREACT Full-text

TITLE: Process for acylation of (hydroxy)-containing aromatic compounds, particularly benzothiophenes, with aromatic hydroxycarboxylic acids in the presence of Lewis acids and halogenosilanes

INVENTOR(S): Bourgeois, Damien

PATENT ASSIGNEE(S): Rhodia Chimie, Fr.

SOURCE: Fr. Demande, 35 pp.

CODEN: FRXXBL

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 1



PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2877341	A1	20060505	FR 2004-11646	20041102
CA 2585714	A1	20060511	CA 2005-2585714	20051028
WO 2006048545	A1	20060511	WO 2005-FR2716	20051028
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM EP 1809617 A1 20070725 EP 2005-815207 20051028 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR IN 2007DN03286 A 20070831 IN 2007-DN3286 20070501 PRIORITY APPLN. INFO.: FR 2004-11646 20041102 WO 2005-FR2716 20051028				

OTHER SOURCE(S): MARPAT 144:450603

AB The invention is related to a process for the acylation of aromatic compds., particularly benzothiophenes I [R4 = alkyl, halogenophenyl, (un)substituted Ph; each R5 = independently H, NO2, alkyl, alkoxy, halo, CF3, etc.; n = 0-3], with aromatic hydroxycarboxylic acids II [each R7 = H or a substituent, especially alkyl, alkoxy, NO2, CN; m < 4], in the presence of a Lewis acid and a halogenosilane to give the ketones III. The advantages include acylation of hydroxy-containing substrates and/or agents without OH group protection, absence of toxic materials and simple procedure. Thus, successive addition of 4-hydroxybenzoic acid, chlorobenzene, methyltrichlorosilane, 2-butyl-5-nitrobenzofuran (IV) and FeCl3 at 23°, and stirring at 40° for 5 h gave 2-butyl-3-(4-hydroxybenzoyl)-5-nitrobenzofuran in 78% selectivity at 95% conversion of IV.

RX(1) OF 1 A + B ==> C



C
YIELD 82%

RX(1) RCT A 99-96-7

STAGE(1)

SOL 108-90-7 PhCl

CON room temperature -> 40 deg C

STAGE(2)

RGT D 7705-08-0 FeCl3, E 75-79-6 MeSiCl3

CON 15 minutes, 40 deg C

STAGE(3)

RCT B 133238-87-6

SOL 108-90-7 PhCl

CON SUBSTAGE(1) 12 minutes, 40 deg C

 SUBSTAGE(2) 3 hours, 40 deg C

 SUBSTAGE(3) 40 deg C -> 30 deg C

STAGE(4)

SOL 64-17-5 EtOH

CON SUBSTAGE(1) 17 minutes, 30 deg C

 SUBSTAGE(2) 10 minutes, 30 deg C

PRO C 141645-16-1

NTE optimization study, optimized on temperature, order and mode of
 addition of reaction participants

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L19 ANSWER 2 OF 6 CASREACT COPYRIGHT 2007 ACS on STN DUPLICATE 2

ACCESSION NUMBER: 143:97254 CASREACT [Full-text](#)

TITLE: Process for preparation de 2-(n-alkyl)-3-(4-hydroxybenzoyl)benzofurans and intermediates by halogenation of carboxybenzofuran derivatives, Friedel-Crafts acylation with alkoxybenzenes and dealkylation

INVENTOR(S): Schouteeten, Alain; Bleger, Francois; Mordacq,


Nizal Chandrakumar 10/584,440

PATENT ASSIGNEE(S): Francoise; Piron, Jerome
 SOURCE: Clariant France, Fr.
 DOCUMENT TYPE: Fr. Demande, 22 pp.
 LANGUAGE: CODEN: FRXXBL
 French Patent
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2864536	A1	20050701	FR 2003-15398	20031224
FR 2864536	B1	20060317		
WO 2005066149	A1	20050721	WO 2004-IB4158	20041215
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RM:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GD, GW, ML, MR, NE, SN, TD, TG			
EP 1699772	A1	20060913	EP 2004-801395	20041215
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS			
CN 1898226	A	20070117	CN 2004-80038285	20041215
JP 2007517012	T	20070628	JP 2006-546365	20041215
NO 2006002936	A	20060922	NO 2006-2936	20060623
IN 2006CN02324	A	20070706	IN 2006-CN2324	20060626
US 2007155831	A1	20070705	US 2006-584440	20061129
PRIORITY APPLN. INFO.:			FR 2003-15398	20031224
			WO 2004-IB4158	20041215

AB The invention is related to the preparation of benzofurans I [R = linear or branched alkyl; R1 = halo, NO2, linear or branched alkyl, alkoxy] and intermediates by halogenation of acids II [R1, R defined as above] in an organic solvent, Friedel-Crafts acylation of alkoxybenzenes of formula C6H5OR2 (III) [R2 = linear or branched alkyl] with acyl halides IV (X = halo) in the presence of a Lewis acid to V [R, R1, R2 defined as above] and its 2-alkoxy isomer, and dealkylation. The invention is also related to the preparation of II by heating VI [R1' = NO2; R4 = linear or branched alkyl] and its ketone tautomer in the presence of an acid catalyst. The advantages include absence of poisoned materials, higher yields and purities. For example, chlorination of 2-(n-butyl)-3-carboxy-5-nitrobenzofuran with SOCl2 in PhCl, acylation of anisole with acyl chloride in the presence of AlCl3, and demethylation over AlCl3 at 60° for 7 h gave a solid containing 99.5% I [R1 = 5-NO2, R = n-Bu] after purification. Heating 3-(1-hydroxypentylidene)-5-nitro-2(3H)-benzofuran in the presence of acetic anhydride/H2SO4 for 2 h gave acid II (m.p. = 207°).

RX(3) OF 14 ...F ==> I



RX(3) RCT F 141627-42-1

STAGE(1)

RGT H 7446-70-0 AlCl₃

SOL 108-90-7 PhCl

CON 7 hours, room temperature -> 60 deg C

STAGE(2)

RGT J 7732-18-5 Water

PRO I 141645-16-1

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L19 ANSWER 3 OF 6 CASREACT COPYRIGHT 2007 ACS on STN DUPLICATE 3

ACCESSION NUMBER: 122:105333 CASREACT [Full-text](#)

TITLE: Regioselectivity in the Alkaline Thiolate Deprotection
of Aryl Methyl Ethers

AUTHOR(S): Dodge, Jeffrey A.; Stocksdales, Mark G.; Fahey, Kennan
J.; Jones, C. David

CORPORATE SOURCE: Lilly Research Laboratories, Eli Lilly and Company,
Indianapolis, IN, 46285, USA

SOURCE: Journal of Organic Chemistry (1995), 60(3), 739-41
CODEN: JOCEAH; ISSN: 0022-3263


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal


LANGUAGE: English

AB The regioselective deprotection of aryl Me ethers using sodium ethanethiolate in DMF was systematically explored. Electronic factors appear to control the observed selectivity, with Me ethers para to electron-withdrawing groups reacting preferentially with the thiol anion. In addition, substituent effects indicate a relationship between the Hammett constant and the efficacy of the reaction, with more electron-poor species providing higher yields of demethylated product. A variety of these substituents (NO₂, CN, acetyl) provide useful yields of deprotected product, thereby adding synthetic utility to this general method.

RX(4) OF 15 I ==> J



I

(4)
→

J

YIELD 75%

RX(4) RCT I 160663-54-7
 RGT C 811-51-8 NaSEt
 PRO J 160663-56-9
 SOL 68-12-2 DMF
 NTE REGIOSELECTIVE

L19 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2007 ACS on SIN
 ACCESSION NUMBER: 2006:815935 CAPLUS Full-text
 145:230520

DOCUMENT NUMBER: 145:230520
 TITLE: Preparation of benzofurans and related derivatives as
 tubulin polymerization inhibitors for treating
 neoplasm and inflammation

INVENTOR(S): Chaplin, Jason Hugh; Gill, Gurmit Singh; Grobelny,
 Damian Wojciech; Flynn, Bernard Luke

PATENT ASSIGNEE(S): Iliad Chemicals Pty Ltd, Australia

SOURCE: PCT Int. Appl., 147pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006084338	A1	20060817	WO 2006-AU192	20060214
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX,				

MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, VZ, VN, YU, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

AU 2006212726 A1 20060817 AU 2006-212726 20060214
CA 2597447 A1 20060817 CA 2006-2597447 20060214
EP 1848704 A1 20071031 EP 2006-704869 20060214


R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR

PRIORITY APPLN. INFO.: US 2005-652668P P 20050214
WO 2006-AU192 W 20060214


OTHER SOURCE(S): MARPAT 145:230520

ED Entered SIN: 17 Aug 2006

GI



I



II

AB Title compds. I [X = O, S, SO, SO₂, Se, SeO, SeO₂, NH and derivs.; R1-R4 = independently H, CO₂H, CN, OH, NO₂, (un)substituted acyl, arylalkoxy, aryl, oxyacylamino, etc.; Y = (un)substituted Ph, phenylcarbonyl, phenoxy, phenylsulfanyl, etc.; Q = (un)substituted heteroaryl, heterocyclyl, heteroarylcarbonyl, etc.; with provisios; and their salts] were prepared as tubulin polymerization inhibitors. Thus, Sonogashira coupling of 2,4-dimethoxy-3-nitroiodobenzene (preparation given) with 4-ethynyl-1-methyl-1H-pyrazole, cyclization with bis(pyridine)iodonium tetrafluoroborate, Stille coupling with trimethyl(3,4,5-trimethoxyphenyl)stannane, and reduction with Zn/AcOH gave benzofuran II. II inhibited tubulin polymerization (IC₅₀ = 1.5±0.1 μM). Selected I inhibited proliferation of MCF-7 breast cancer cells and activated HUVEC cells. I are useful for treating neoplasm and inflammation.

CC 27-7 (Heterocyclic Compounds (One Hetero Atom))

Section cross-reference(s): 1

IT 905751-61-3P, [2-(1-Benzyl-1H-pyrazol-4-yl)-6-methoxybenzofuran-3-yl](3,4,5-trimethoxyphenyl)methanone 905751-63-5P, [7-Hydroxy-6-methoxy-2-(1H-pyrazol-4-yl)benzofuran-3-yl](3,4,5-trimethoxyphenyl)methanone 905751-68-0P, 2-(1-Methylpyrazol-4-yl)-3-(3,4,5-trimethoxybenzoyl)-6-

methoxy-7-hydroxybenzofuran 905751-70-4P, 2-(1-Methylpyrazol-4-yl)-3-(3,5-dimethoxybenzoyl)-6-methoxy-7-hydroxybenzofuran 905751-75-9P
 905751-76-0P, 2-[4-(6-Methoxy-3-(3,4,5-trimethoxybenzoyl)benzo[b]furan-2-yl)pyrazol-1-yl]acetamide 905751-77-1P, [6-Methoxy-2-[1-(4-methoxyphenyl)-1H-pyrazol-4-yl]benzofuran-3-yl](3,4,5-trimethoxyphenyl)methanone 905751-78-2P, [2-[1-(2-Dimethylaminoethyl)-1H-pyrazol-4-yl]-6-methoxybenzofuran-3-yl](3,4,5-trimethoxyphenyl)methanone 905751-79-3P, 2-[4-[7-Hydroxy-6-methoxy-3-(3,4,5-trimethoxybenzoyl)benzofuran-2-yl]pyrazol-1-yl]acetamide 905751-81-7P, [2-(1-Methyl-1H-1,2,4-triazol-4-yl)-6-methoxybenzofuran-3-yl](3,4,5-trimethoxyphenyl)methanone 905751-82-8P 905751-85-1P 905751-88-4P 905751-91-9P 905751-95-3P 905751-97-5P, (2S)-2-Amino-3-hydroxy-N-[6-methoxy-2-(1-methyl-1H-pyrazol-4-yl)-3-(3,4,5-trimethoxybenzoyl)benzofuran-7-yl]propanamide Hydrochloride 905752-01-4P 905752-03-6P, [6-Methoxy-2-(1-methyl-1H-pyrazol-4-yl)-1H-indol-3-yl](3,4,5-trimethoxyphenyl)methanone 905752-09-2P, [6-Methoxy-7-nitro-2-(1-methyl-1H-pyrazol-4-yl)benzofuran-3-yl](3,4,5-trimethoxyphenyl)methanone 905752-10-5P, 7-Amino-6-methoxy-2-(1-methyl-1H-pyrazol-4-yl)-3-[(3,4,5-trimethoxyphenyl)thio]benzo[b]furan 905752-12-7P, [7-Fluoro-6-methoxy-2-(1-methyl-1H-pyrazol-4-yl)benzofuran-3-yl](3,4,5-trimethoxyphenyl)methanone 905752-16-1P, 2-[4-[7-Fluoro-6-methoxy-3-(3,4,5-trimethoxybenzoyl)benzofuran-2-yl]-1H-pyrazol-1-yl]acetamide 905752-18-3P 905752-22-9P 905752-40-1P, 2-(6-Methoxypyridin-3-yl)-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzofuran 905752-42-3P, 2-(Thiophen-3-yl)-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-43-4P, 2-(3,5-Dimethylisoxazol-4-yl)-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-44-5P, 2-(1-Isobutylpyrazol-4-yl)-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-45-6P, 2-[5-(Formyl)thiophen-2-yl]-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-46-7P, 2-(1-1,2,4-triazol-4-yl)-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-47-8P, 2-(1,2,3-Triazol-1-yl)-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-48-9P, 2-(1-Pyrazolyl)-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-49-0P, 2-(1,2,4-Triazol-1-yl)-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-50-3P, 2-(1-Pyrrolyl)-7-hydroxy-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-51-4P, 2-(4-Methylpiperazino)-3-(3,4,5-trimethoxybenzoyl)-6-methoxybenzo[b]furan 905752-52-5P, 2-(2-Furyl)-6-methoxy-3-(3,4,5-trimethoxybenzoyl)benzo[b]furan 905752-54-7P, 7-Hydroxy-6-methoxy-2-(2H-tetrazol-5-yl)-3-(3,4,5-trimethoxybenzoyl)benzo[b]furan 905752-56-9P, 7-Hydroxy-6-methoxy-2-(2H-[1,2,3]triazol-4-yl)-3-(3,4,5-trimethoxybenzoyl)benzo[b]furan 905752-58-1P 905752-59-2P 905752-62-7P, [4-[6-Methoxy-3-(3,4,5-trimethoxybenzoyl)benzo[b]furan-2-yl]pyrazol-1-yl]acetic acid 905752-63-8P, (2S)-2-Amino-3-hydroxy-N-[6-methoxy-2-(1-methyl-1H-pyrazol-4-yl)-3-(3,4,5-trimethoxybenzoyl)benzofuran-7-yl]propanamide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation);
 USES (Uses)

(drug candidate; preparation of benzofurans and related derivs. as tubulin polymerization inhibitors for treating neoplasm and inflammation)

IT 4371-79-3, Carbon diiodide 7726-95-6, Bromine, reactions 7789-33-5, Iodine bromide (IBr)

RL: RCT (Reactant); RGT (Reagent); RACT (Reactant or reagent) (biologisation agent; preparation of benzofurans and related derivs. as tubulin polymerization inhibitors for treating neoplasm and inflammation)

IT 905752-18-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU


Nizal Chandrakumar 10/584,440

(Therapeutic use); BIOL (Biological study); PREP (Preparation);
USES (Uses)

(drug candidate; preparation of benzofurans and related derivs. as
tubulin polymerization inhibitors for treating neoplasm and inflammation)

RN 905752-18-3 CAPLUS

CN Methanone, (4-hydroxy-3,5-dimethoxyphenyl) [7-hydroxy-6-methoxy-2-(1-methyl-
1H-pyrazol-4-yl)-3-benzofuranyl]- (CA INDEX NAME)



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L19 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 2005:1329213 CAPLUS Full-text

DOCUMENT NUMBER: 144:51437

TITLE: Xanthine oxidase inhibitor, 6-hydroxybenzobromarone,
and process for the preparation thereof

INVENTOR(S): Endou, Hitoshi; Oikawa, Toshihiro

PATENT ASSIGNEE(S): Torii Pharmaceutical Co., Ltd., Japan; Human Cell
Systems, Inc.

SOURCE: PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005121112	A1	20051222	WO 2005-JP10671	20050610
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, VZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1767531	A1	20070328	EP 2005-748953	20050610
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR			
US 2007185195	A1	20070809	US 2006-628918	20061206
PRIORITY APPLN. INFO.:			JP 2004-172456	A 20040610
			WO 2005-JP10671	W 20050610

ED Entered STN: 22 Dec 2005

- AB Process for the preparation of 6-hydroxybenzobromarone (I) was provided. Thus, 2-ethyl-3-(p-hydroxybenzoyl)-6-methoxybenzofuran (6.75 mmol) was reacted with NBS (67.5 mmol) in CH₂Cl₂ at room temperature for 15 h to give 6-methoxybenzobromarone in 55% yield. Treatment of 6-methoxybenzobromarone (3.74 mmol) with AlCl₃ (17.6 mmol) and ethanethiol (7 mL) in CH₂Cl₂ (35 mL) at ice-bath temperature for 10 min followed by acid work-up and silica gel purification afforded 6-hydroxybenzobromarone in 63% yield. In xanthine oxidase inhibition assays, the IC₅₀ value of compound I was 68 μM. Compound I is claimed useful for the treatment of hyperuricemia, gout, etc.
- IC ICM C07D307-80
- ICS A61K031-343; A61P013-02; A61P019-02; A61P019-06; A61P043-00
- CC 27-7 (Heterocyclic Compounds (One Hetero Atom))
- Section cross-reference(s): 1
- IT Dealkylation
(preparation of 6-hydroxybenzobromarone via deprotection of 6-methoxybenzobromarone)
- IT 871493-08-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP
{Preparation}; RACT (Reactant or reagent)
(bromination of 2-ethyl-3-(p-hydroxybenzoyl)-6-methoxybenzofuran using N-bromosuccinimide)
- IT 871493-07-1P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP
{Preparation}; RACT (Reactant or reagent)
(demethylation of 3-(p-anisoyl)-2-ethyl-6-methoxybenzofuran using ethanethiol sodium salt)
- IT 871493-08-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP
{Preparation}; RACT (Reactant or reagent)
(bromination of 2-ethyl-3-(p-hydroxybenzoyl)-6-methoxybenzofuran using N-bromosuccinimide)
- RN 871493-08-2 CAPLUS
- CN Methanone, (3,5-dibromo-4-hydroxyphenyl) (2-ethyl-6-methoxy-3-benzofuranyl)- (CA INDEX NAME)



- IT 871493-07-1P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP
{Preparation}; RACT (Reactant or reagent)
(demethylation of 3-(p-anisoyl)-2-ethyl-6-methoxybenzofuran using ethanethiol sodium salt)
- RN 871493-07-1 CAPLUS
- CN Methanone, (2-ethyl-6-methoxy-3-benzofuranyl) (4-hydroxyphenyl)- (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L19 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 1961:76079 CAPLUS Full-text

DOCUMENT NUMBER: 55:76079

ORIGINAL REFERENCE NO.: 55:14420e-i,14421a-e

TITLE: Study of benzofuran. V. Structure of the diketones obtained from the acylation of 2-ethyl-3-acylbenzofurans

AUTHOR(S): Bisagni, Emile; Royer, Rene

CORPORATE SOURCE: Inst. radium, Paris

SOURCE: Bulletin de la Societe Chimique de France (1960) 1968-76

CODEN: BSCFAS; ISSN: 0037-8968

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

ED Entered STN: 22 Apr 2001

AB cf. CA 54, 24632a; 55, 505b. 2-Ethyl-3-benzoyl- and 2-ethyl-3-anisoylbenzofurans have been acetylated under Friedel-Crafts conditions with excess AlCl₃. Substitution occurred first at the 6-position, next at the 5-position. 2-Acylbenzofurans were not acetylated under the same conditions. To a solution of 1 mole 2-ethyl-3-benzoylbenzofuran (I) and 2 moles AcCl in 700 cc. CS₂ was gradually added 2.5 moles AlCl₃. The mixture was kept 24 hrs., then decomposed and purified to yield 4.5% I and a mixture, b. 240-60°, which on fractional crystallization from EtOH gave 40% 2-ethyl-3-benzoyl-6-acetylbenzofuran (II), m. 118.5° and 28% 2-ethyl-3-benzoyl-5-acetylbenzofuran (III), m. 68° (ligroine). Similarly, 2-ethyl-3-(4-methoxybenzoyl)benzofuran (IV) with AcCl gave 45% 2-ethyl-3-(4-methoxybenzoyl)-6-acetylbenzofuran (V), m. 118.5-19°, and 7% 2-ethyl-3-(4-methoxybenzoyl)-5-acetylbenzofuran (VI), m. 100-1° (EtOH, then ligroine-C₆H₆). V was demethylated by refluxing 20 min. with pyridine-HCl to 2-ethyl-3-(4-hydroxybenzoyl)-6-acetylbenzofuran, m. 214-15° (EtOH, or C₆H₆-ligroine). VI did not respond to similar treatment. NaOH degradation of II gave 4,2-Ac(HO)C₆H₃CH₂COPh (VII), m. 212-13°, BzOH, and 4,2-Ac(HO)C₆H₃CH₂COEt (VIII). Treatment of III with NaOH gave BzOH, 5,2-Ac(HO)C₆H₃CH₂COPh (IX), m. 179°, and 5,2-Ac(HO)C₆H₃CH₂COEt (X). Similarly, V with NaOH gave 4,2-Ac(HO)C₆H₃CH₂CO₂C₆H₄OMe-p (XI), m. 211°, anisic acid, and VIII. NaOH degradation of VI yielded 5,2-Ac(HO)C₆H₃CH₂CO₂C₆H₄OMe-p (XII), m. 165-7°, anisic acid, and X. The mixture of VII and VIII obtained from the NaOH degradation of II was methylated with MeI to yield 31% 4,2-Ac(MeO)C₆H₃CH₂COEt (XIII), b₁₇ 201-4°, n₂₂ 1.5390, and 25.5% 4,2-Ac(MeO)C₆H₃CH₂COPh (XIV), b₁₇ 247-8°, m. 66° (ligroine). Methylation of the mixture of VIII and XI gave 11% XIII and 51.1% 4,2-Ac(MeO)C₆H₃CH₂CO₂C₆H₄OMe-p (XV), b₁₆ 275-8°, m. 69-70° (ligroine-20% cyclohexane). Heating VIII in EtOH saturated with HCl gave 90% 2-ethyl-6-acetylbenzofuran (XVI), b₁₂ 163-5°, n_{20.5} 1.5845, m. 20-2°; oxime m. 93.5° (dilute Et₂O or ligroine). NaOB treatment of XVI gave 33% 2-ethyl-6-benzofurancarboxylic acid, m. 171-2°. XVI was reduced by N₂H₄ in (CH₂OH)₂ to 2,6-diethylbenzofuran, b₁₅ 126.5°, n_{22.5} 1.5415. In the same way, VII, heated in EtOH saturated with HCl gave 2-phenyl-6-acetylbenzofuran, m. 103-4° which was reduced by N₂H₄ to 2-phenyl-6-ethylbenzofuran, m. 52-3° (EtOH). XI heated in EtOH saturated with HCl gave 80% 2-(4-methoxyphenyl)-6-acetylbenzofuran, m. 147° (EtOH-C₆H₆), which was demethylated to 2-(4-hydroxyphenyl)-6-acetylbenzofuran, m. 228° (EtOH or C₆H₆) and reduced by N₂H₄ to 2-(4-hydroxyphenyl)-6-ethylbenzofuran, m. 170-1° (dilute EtOH), b₁₄ 244-7°. X heated in EtOH saturated with HCl gave 2-ethyl-5-acetylbenzofuran, b₂₃ 179-80°, m. 44-5°; oxime m. 83°. The latter, treated with NaOB gave 2-ethyl-5-benzofurancarboxylic acid, m. 165° (dilute EtOH). Cyclization of IX yielded 2-phenyl-5-acetylbenzofuran, m. 160° (EtOH), which was reduced by N₂H₄ in

(CH₂OH)₂ to 2-phenyl-5-ethylbenzofuran, m. 76° (EtOH), b22 212-14°. XII was also cyclized (EtOH-HCl) to give 68% 2-(4-methoxyphenyl)-5-acetylbenzofuran, m. 170° (EtOH-C₆H₆), which was simultaneously reduced and demethylated by N₂H₄ to 2-(4-hydroxyphenyl)-5-ethylbenzofuran, m. 186° (dilute EtOH or C₆H₆). 5-Ethylsalicylaldehyde, b16 115-16°, n_D1.5 1.5545, was treated with ClCH₂-CO₂Me and KOH in EtOH to yield 5% 2-acetyl-5-ethylbenzofuran, b15 163-4°, m. 33-4° (EtOH), which was reduced to 2,5-diethylbenzofuran (XVII), b14 122-4°, n_D1.5430. XVII was benzoylated (ClCOPh, SnCl₄, C₆H₆) in 67% yield to give 2,5-diethyl-3-benzoylbenzofuran (XVIII), b13 220-2°, n_D20 1.5995. Anisoylation of XVII gave 2,5-diethyl-3-(4-methoxybenzoyl)benzofuran (XIX), b12 247°, m. 32-3° (EtOH). The latter was demethylated to 2,5-diethyl-3-(4-hydroxybenzoyl)benzofuran, b4 272-3°, m. 135-6° (C₆H₆). NaOH degradation of XVIII followed by HCl-EtOH recyclization gave 43% PhCO₂H, 22.5% XVII, and 40% 2-phenyl-5-ethylbenzofuran. Similar treatment of XIX gave 29.5% anisic acid, 2-(4-methoxyphenyl)-5-ethylbenzofuran, m. 135°, 5,2-Et (HO)C₆H₃CH₂CO₂Me-p, m. 118°. Starting with 3-ethylsalicylaldehyde, b28 117-18°, a similar series of reactions was carried out giving 2-acetyl-7-ethylbenzofuran, b19 159-61°, m. 54.5° (EtOH), 2,7-diethylbenzofuran (XX), b20 124-5°, n_D22 1.5410, and 2,7-diethyl-3-benzoylbenzofuran, b15 228-31°, n_D21.5 1.6038. NaOH degradation of the latter gave BzOH, XX, and 2-phenyl-7-ethylbenzofuran, b20 220-3°, n_D24 1.6210.

CC 10G (Organic Chemistry: Heterocyclic Compounds)

IT Acylation

(of 3-acyl-2-ethylbenzofurans, structure of diketones from)

IT Ketones


(structure of di-, from acylation of 3-acyl-2-ethylbenzofurans)

IT 3131-63-3, Benzofuran, 2-ethyl-


(3-acyl derivs., diketones from acylation of)

IT 5896-26-4P, Ketone, 2-ethyl-6-benzofuranyl methyl 5896-49-1P, Ketone, 2,6-diethyl- 27408-42-0P, Ketone, 2-ethyl-6-benzofuranyl methyl, oxime 28089-83-0P, Ketone, methyl 2-phenyl-6-benzofuranyl 59664-03-8P, Ketone, 7-ethyl-2-benzofuranyl methyl 91495-47-5P, Benzofuran, 2,5-diethyl- 93021-68-2P, Benzofuran, 5-ethyl-2-phenyl- 94066-54-3P, 2,4'''-Biacetophenone, 3'''-hydroxy- 94302-86-0P, Ketone, 2,5-diethyl-3-benzofuranyl p-hydroxyphenyl 95485-40-8P, Ketone, 2-ethyl-5-benzofuranyl methyl 100612-37-1P, Acetophenone, 3'-methoxy-4'-(2-oxobutyl)- 101278-17-5P, Ketone, 2-(p-hydroxyphenyl)-6-benzofuranyl methyl 101594-95-0P, Acetophenone, 2-(5-ethyl-2-hydroxyphenyl)-4'-methoxy- 101596-59-2P, Benzofuran, 5-ethyl-2-(p-methoxyphenyl)- 101894-27-3P, Benzofuran, 6-acetyl-2-ethyl-3-p-hydroxybenzoyl- 101894-27-3P, Phenol, p-(6-acetyl-2-ethyl-3-benzofuranylcarbonyl)- 102158-98-5P, Ketone, 2,5-diethyl-3-benzofuranyl p-methoxyphenyl 103152-25-6P, 2,4'''-Biacetophenone, 3'''-methoxy- 103988-06-3P, 5-Benzofurancarboxylic acid, 2-ethyl- 105207-89-4P, Acetophenone, 4'-hydroxy-3'-(2-oxobutyl)- 105208-20-6P, Acetophenone, 3'-hydroxy-4'-(2-oxobutyl)- 105909-84-0P, Ketone, 2-ethyl-5-benzofuranyl methyl, oxime 106989-39-3P, Ketone, 5-ethyl-2-benzofuranyl methyl 108838-38-6P, 2,3'''-Biacetophenone, 4'''-hydroxy- 108840-63-7P, Benzofuran, 6-ethyl-2-phenyl- 108840-64-8P, Benzofuran, 7-ethyl-2-phenyl- 108840-82-0P, Phenol, p-6-ethyl-2-benzofuranyl- 108842-68-8P, Phenol, p-5-ethyl-2-benzofuranyl- 108980-53-6P, Ketone, 2-(p-methoxyphenyl)-6-benzofuranyl methyl 108983-46-6P, Ketone, 2-(p-methoxyphenyl)-5-benzofuranyl methyl 109155-31-9P, 2,4'''-Biacetophenone, 3'''-hydroxy-4'-methoxy- 109156-66-3P, 2,3'''-Biacetophenone, 4'''-hydroxy-3'-methoxy- 109395-02-0P, 2,4'''-Biacetophenone, 3'''', 4'-dimethoxy- 109614-26-8P, Benzofuran, 6-acetyl-3-benzoyl-2-ethyl- 109614-90-6P, Benzofuran, 5-acetyl-3-benzoyl-2-ethyl- 109688-24-6P, Ketone, 2,7-diethyl-3-

benzofuranyl phenyl 109690-79-1P, Ketone, 2,5-diethyl-3-benzofuranyl phenyl 109893-46-1P, 1-Propanone, 1-[5(or 6)-acetyl-2-ethyl-3-benzofuranyl]- 109936-61-0P, Benzofuran, 6-acetyl-3-p-anisoyl-2-ethyl-109938-44-5P, Benzofuran, 5-acetyl-3-p-anisoyl-2-ethyl- 121045-41-8P, Ketone, methyl 2-phenyl-5-benzofuranyl 857020-75-8P, 6-Benzofurancarboxylic acid, 2-ethyl- 857021-42-2P, Benzofuran, 2,7-diethyl-
 RL: PREP (Preparation)
 (preparation of)
 IT 94302-86-0P, Ketone, 2,5-diethyl-3-benzofuranyl p-hydroxyphenyl 101894-27-3P, Benzofuran, 6-acetyl-2-ethyl-3-p-hydroxybenzoyl-
 RL: PREP (Preparation)
 (preparation of)
 RN 94302-86-0 CAPLUS
 CN Ketone, 2,5-diethyl-3-benzofuranyl p-hydroxyphenyl (6CI, 7CI) (CA INDEX NAME)



RN 101894-27-3 CAPLUS
 CN Phenol, p-(6-acetyl-2-ethyl-3-benzofuranylcarbonyl)- (6CI) (CA INDEX NAME)



L27 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:569050 CAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:97254
 TITLE: Process for preparation de 2-(n-alkyl)-3-(4-hydroxybenzoyl)benzofurans and intermediates by halogenation of carboxybenzofuran derivatives, Friedel-Crafts acylation with alkoxybenzenes and dealkylation
 INVENTOR(S): Schouteeten, Alain; Bieger, Francois ; Mordacq, Francoise; Firon, Jerome
 PATENT ASSIGNEE(S): Clariant France, Fr.
 SOURCE: Fr. Demande, 22 pp.
 CODEN: FRXXBL
 DOCUMENT TYPE: Patent
 LANGUAGE: French
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

Nizal Chandrakumar 10/584,440

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2864536	A1	20050701	FR 2003-15398	20031224
FR 2864536	B1	20060317		
WO 2005066149	A1	20050721	WO 2004-IB4158	20041215
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1699772	A1	20060913	EP 2004-801395	20041215
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS				
CN 1898226	A	20070117	CN 2004-80038285	20041215
JP 2007157012	T	20070628	JP 2006-546365	20041215
NO 2006002936	A	20060922	NO 2006-2936	20060623
IN 2006CN02324	A	20070706	IN 2006-CN2324	20060626
US 2007155831	A1	20070705	US 2006-584440	20061129
PRIORITY APPLN. INFO.:			FR 2003-15398	A 20031224
			WO 2004-IB4158	W 20041215
OTHER SOURCE(S): CASREACT 143:97254				
ED Entered STN: 01 Jul 2005				
GI				

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

- AB The invention is related to the preparation of benzofurans I [R = linear or branched alkyl; R1 = halo, NO2, linear or branched alkyl, alkoxy] and intermediates by halogenation of acids II [R1, R defined as above] in an organic solvent, Friedel-Crafts acylation of alkoxybenzenes of formula C6H5OR2 (III) [R2 = linear or branched alkyl] with acyl halides IV (X = halo) in the presence of a Lewis acid to V [R, R1, R2 defined as above] and its 2-alkoxy isomer, and dealkylation. The invention is also related to the preparation of II by heating VI [R1' = NO2; R4 = linear or branched alkyl] and its ketone tautomer in the presence of an acid catalyst. The advantages include absence of poisoned materials, higher yields and purities. For example, chlorination of 2-(n-butyl)-3-carboxy-5-nitrobenzofuran with SOCl2 in PhCl, acylation of anisole with acyl chloride in the presence of AlCl3, and demethylation over AlCl3 at 60° for 7 h gave a solid containing 99.5% I [R1 = 5-NO2, R = n-Bu] after purification. Heating 3-(1-hydroxypentylidene)-5-nitro-2(3H)-benzofuran in the presence of acetic anhydride/H2SO4 for 2 h gave acid II (m.p. = 207°).
- IC ICM C07D307-80
- CC 27-7 (Heterocyclic Compounds (One Hetero Atom))
Section cross-reference(s): 45
- IT 256758-95-99P, 2-(n-Butyl)-3-(2-hydroxybenzoyl)-5-nitrobenzofuran
RL: BYP (Byproduct); IMF (Industrial manufacture); RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)
(process for preparation de 2-(n-alkyl)-3-(4-hydroxybenzoyl)benzofurans and intermediates by halogenation of the corresponding carboxybenzofurans, Friedel-Crafts acylation with alkoxybenzenes and dealkylation)
- IT i41645-i6-1P, 2-(n-Butyl)-3-(4-hydroxybenzoyl)-5-nitrobenzofuran

Nizal Chandrakumar 10/584,440

RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP
(Preparation)

(product; process for preparation de 2-(n-alkyl)-3-(4-
hydroxybenzoyl)benzofurans and intermediates by halogenation of the
corresponding carboxybenzofurans, Friedel-Crafts acylation with
alkoxybenzenes and dealkylation)

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>